

Outpatient[†] management of skin and soft tissue infections in the era of community-associated MRSA[‡]

Patient presents with signs/symptoms of skin infection:

- Redness
- Swelling
- Warmth
- Pain/tenderness
- Complaint of “spider bite”

YES

Is the lesion purulent (i.e., are any of the following signs present)?

- Fluctuance—palpable fluid-filled cavity, movable, compressible
- Yellow or white center
- Central point or “head”
- Draining pus
- Possible to aspirate pus with needle and syringe

NO

Possible cellulitis without abscess:

- Provide antimicrobial therapy with coverage for *Streptococcus* spp. and/or other suspected pathogens
- Maintain close follow-up
- Consider adding coverage for MRSA (if not provided initially), if patient does not respond

YES

1. Drain the lesion
2. Send wound drainage for culture and susceptibility testing
3. Advise patient on wound care and hygiene
4. Discuss follow-up plan with patient

If systemic symptoms, severe local symptoms, immunosuppression, or failure to respond to I&D

Consider antimicrobial therapy with coverage for MRSA in addition to I&D

(See reverse for options)



The use of the CDC logo on this material does not imply endorsement of AMA products/services or activities promoted or sponsored by the AMA.

Options for empiric outpatient antimicrobial treatment of SSTIs when MRSA is a consideration*

Drug name	Considerations	Precautions**
Clindamycin	<ul style="list-style-type: none"> ■ FDA-approved to treat serious infections due to <i>S. aureus</i> ■ D-zone test should be performed to identify inducible clindamycin resistance in erythromycin-resistant isolates 	<ul style="list-style-type: none"> ■ <i>Clostridium difficile</i>-associated disease, while uncommon, may occur more frequently in association with clindamycin compared to other agents.
Tetracyclines <ul style="list-style-type: none"> ■ Doxycycline ■ Minocycline 	<ul style="list-style-type: none"> ■ Doxycycline is FDA-approved to treat <i>S. aureus</i> skin infections. 	<ul style="list-style-type: none"> ■ Not recommended during pregnancy. ■ Not recommended for children under the age of 8. ■ Activity against group A streptococcus, a common cause of cellulitis, unknown.
Trimethoprim-Sulfamethoxazole	<ul style="list-style-type: none"> ■ Not FDA-approved to treat any staphylococcal infection 	<ul style="list-style-type: none"> ■ May not provide coverage for group A streptococcus, a common cause of cellulitis ■ Not recommended for women in the third trimester of pregnancy. ■ Not recommended for infants less than 2 months.
Rifampin	<ul style="list-style-type: none"> ■ Use only in combination with other agents. 	<ul style="list-style-type: none"> ■ Drug-drug interactions are common.
Linezolid	<ul style="list-style-type: none"> ■ Consultation with an infectious disease specialist is suggested. ■ FDA-approved to treat complicated skin infections in adults, including those caused by MRSA. 	<ul style="list-style-type: none"> ■ Has been associated with myelosuppression, neuropathy and lactic acidosis during prolonged therapy.

- MRSA is resistant to all currently available beta-lactam agents (penicillins and cephalosporins)
- Fluoroquinolones (e.g., ciprofloxacin, levofloxacin) and macrolides (erythromycin, clarithromycin, azithromycin) are not optimal for treatment of MRSA SSTIs because resistance is common or may develop rapidly.

* Data from controlled clinical trials are needed to establish the comparative efficacy of these agents in treating MRSA SSTIs. Patients with signs and symptoms of severe illness should be treated as inpatients.

** Consult product labeling for a complete list of potential adverse effects associated with each agent.

Role of decolonization

Regimens intended to eliminate MRSA colonization should not be used in patients with active infections. Decolonization regimens may have a role in preventing recurrent infections, but more data are needed to establish their efficacy and to identify optimal regimens for use in community settings. *After treating active infections and reinforcing hygiene and appropriate wound care*, consider consultation with an infectious disease specialist regarding use of decolonization when there are recurrent infections in an individual patient or members of a household.